

Care of young palms

Sufficient attention will have to be paid to the young palms in the early years of growth. The field planted seedlings should be shaded and irrigated properly during the summer months. Irrigation with 45 liters of water once in 4 days has been found to be satisfactory in all soil types. Provision of proper drainage is also equally important in areas subject to water logging. The pits should be cleared of weeds periodically. Soil washed down by the rains and covering the collar of the seedlings should also be removed. The pits should be widened every year before the application of manure. The pits should be gradually filled up as the seedlings grow. The palms should be frequently examined for any insect or fungal attack and necessary remedial measures should be taken up promptly. Regular manuring right from the first year of planting is essential for good vegetative growth, early flowering and bearing and high yield of coconut palms.

Soil and moisture conservation

Proper soil and moisture conservation practices are vital for better performance of coconut palms; especially in sloppy and undulating terrains.

Regular ploughing or digging of the interspaces of coconut palms twice in a year helps in moisture conservation.

The coconut basins can be mulched with coir dust, coconut husks, green leaves, dried leaves, organic wastes, and dried coconut leaves. Mulching should be done before the end of monsoon and before the top soil dries up.

Application of sufficient quantity of organic manure by way of cattle manure, farmyard manure, compost or green leaves improves the soil characteristics and provides nutrients to coconut palms.

Coconut husk incorporation in the soil helps to absorb and retain large quantities of

water for use by the coconut palms. Husk burial can be done in coconut basins or in the interspace. The husk can be buried either in linear trenches taken 3 m away from the trunk between rows of palms or in circular trenches taken around the palm at a distance of 2 m from the trunk. The trenches may be dug with 50 cm width and 50 cm depth. The husks are to be arranged in layers with concave surface facing upwards and covered with soil.

Measures such as contour binding, terracing etc. can be taken up in sloppy lands for soil and moisture conservation.

Harvesting

For matured nuts, meant for fresh consumption and for copra making, 11-12 month old fruits are to be harvested. For tender coconuts, the fruits are harvested at 5 or 6 or 7 months stage with varied levels of tender and tender kernel maturity. Usually, the mature nuts are harvested 6 to 10 times in a year where the production is only for mature nut production. In well maintained and high yielding gardens, bunches are produced regularly, and harvesting is done once in a month.

Ownership & distribution rights © 2022 ICAR-CIARI, Port Blair



For detailed information please contact

DIRECTOR

ICAR– Central Island Agricultural Research Institute

Port Blair—744 105

Andaman and Nicobar Islands, India

Web: <http://icar-ciari.res.in>

Dweep Haritha - Promising dwarf coconut variety



B. Augustine Jerard
V. Damodaran
I. Jaisankar
K. Abirami
and
E. B. Chakurkar



Division of Horticulture and Forestry

ICAR– Central Island Agricultural Research Institute

Port Blair—744 105

Andaman and Nicobar Islands, India

2022



Dweep Haritha

The coconut palm is known to be the most important of all cultivated palms in the world and is the most extensively grown and used nut in the world. Coconut palm is the predominantly grown crop in the Islands used for varied purposes, is popularly called '*Kalpavriksha*' as all parts of the palm are useful. The coconut fruits provide food, edible oil, sweet water and milk for consumption and industrial uses.

Coconut is the predominant crop of Andaman & Nicobar Islands and the crop is closely associated with the socio-culture of island communities. Coconut is considered as one of the remunerative crops of these islands and the main economy of the Islands directly depend on the fortunes of the crop. Traditionally, tall coconut cultivars of these islands are used for copra making and now for many other kernel-based products such as desiccated coconut, VCO. The tall and dwarf cultivars are used for neera extortion in Nicobar Islands. The native dwarf cultivars of these islands which include Andaman Green Dwarf, Andaman Yellow Dwarf and Andaman Orange Dwarf are popularly used for tender coconut purpose and conserved at ICAR-CIARI, Port Blair. Selections from these conserved dwarf accessions are recommended as better varieties for higher tender coconut yield, dwarfness, ornamental value and earliness.

DWEEP HARITHA

It is a selection from Andaman Green Dwarf accession evaluated over 25 years. ICAR-CIARI has identified and selected this high yielding dwarf selection from Andaman Green Dwarf coconut population and released as Dweep Haritha, for promotion in the island areas suitable for tender coconut purpose. The palms are dwarf with slender stem, produces dark green leaves, petiole, rachis and fruits.



The bunches are huge, fruits are round, medium sized, dehusked fruits are round with average weight of 370g. The palms are regular bearers, shorter drooping leaves, not very strong attachment of leaves with the stem when compared to yellow and orange dwarfs considering the observation of bunch buckling. Owing to the good tender nut water quantity (over 350ml), nut yield (over 150 fruits per palm per year), dwarfness and regular production of bunches, it could be a preferred variety to be grown under home gardens. The cultivar has not by any major pests under field conditions but observed to be susceptible to spiraling whitefly when compared to other tall varieties. The palms give better performance with summer irrigation or else the bunch production may get affected.



Planting material and selection of seedlings

The seedlings are selected based on uniform and vigorous growth in terms of leaf production and early leaf splitting. The vigorous seedlings which are one year old, having

minimum of six leaves and girth of over 10 cm at the collar should be selected for planting.

Selection of the site

Soil with a minimum depth of 1.2 meters and fairly good water holding capacity is preferred for coconut cultivation. Shallow soils with underlying hard rock, low lying areas subject to water stagnation and clayey soils are to be avoided.

Preparation of land and planting

Preparation of land for planting depends to a large extent on soil type and environmental factors. If the land is uneven and full of shrubs, the shrubs have to be cleared and land leveled before taking pits. The depth of pits will depend upon the type of soil. In laterite soil with rocky substratum, deeper and wider pits, 1.2 x 1.2 x 1.2 m, may be dug and filled up with loose soil, powdered cow dung and ash up to a depth of 60 cm before planting. In loamy soils with low water table, planting in pit size of 1 x 1 x 1 m filled with top soil to height of 50 cm is recommended. Two layers of coconut husk can be arranged at the bottom of the pit before filling up the soil. This will help in conserving the soil moisture.

Time of planting

In well drained soils where water stagnation is not a problem, seedlings can be transplanted with the beginning of rainy season. If irrigation facilities are available, it is advisable to take up planting at least a month before the rainy season so that the seedlings get well established before the onset of heavy rains.

Spacing

A spacing of 7.5 x 7.5 m is to be followed. This will accommodate 177 palms per ha under the square system of planting. If the triangular system is adopted, an additional 20 to 25 palms can be planted. Wider spacing of 10 m x 10 m provides ample opportunity to accommodate a number of perennial and annual crops in the interspaces.